



- disease: two-year follow-up of the FRISC-II invasive study. *J Am Coll Cardiol* 2002;**40**:1902–14.
- 13 Fox KAA, Poole-Wilson P, Clayton TC, *et al*. 5-year outcome of an interventional strategy in non-ST-elevation acute coronary syndrome: the British Heart Foundation RITA 3 randomized trial. *Lancet* 2005;**366**:914–20.
 - 14 Smith SC Jr, Dove JT, Jacobs AK, *et al*. ACC/AHA guidelines for percutaneous coronary intervention (revision of the 1993 PTCA guidelines)-executive summary: a report of the American College of Cardiology/American Heart Association task force on practice guidelines (Committee to revise the 1993 guidelines for percutaneous transluminal coronary angioplasty) endorsed by the Society for Cardiac Angiography and Interventions. *Circulation* 2001;**103**:3019–41.
 - 15 Braunwald E, Antman EM, Beasley JW, *et al*. ACC/AHA guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction—2002: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Unstable Angina). *Circulation* 2002;**106**:1893–900.
 - 16 Van de Werf F, Gore JM, Avezum A, *et al*. Access to catheterisation facilities in patients admitted with acute coronary syndrome: multinational registry study. *BMJ* 2005;**330**:441–7.
 - 17 Yusuf S, Flather M, Pogue J, *et al*. Variations between countries in invasive cardiac procedures and outcomes in patients with suspected unstable angina or myocardial infarction without initial ST elevation. *Lancet* 1998;**352**:507–14.
 - 18 Ryan JS, Peterson ED, Chen AY, *et al*. Optimal timing of intervention in non-ST-segment elevation acute coronary syndromes. Insights from the CRUSADE (Can rapid risk stratification of unstable angina patients suppress adverse outcomes with early implementation of the ACC/AHA guidelines) registry. *Circulation* 2004;**112**:3049–57.
 - 19 Bavry A, Kumbhani DJ, Quiroz R, *et al*. Invasive therapy along with glycoprotein IIb/IIIa inhibitors improves survival in non-ST-segment elevation acute coronary syndromes: a meta-analysis and review of the literature. *Am J Cardiol* 2004;**93**:830–5.
 - 20 Mehta SR, Cannon CP, Fox KAA, *et al*. Routine vs selective invasive strategies in patients with acute coronary syndromes: a collaborative meta-analysis of randomized trials. *JAMA* 2005;**293**:2908–17.
 - 21 Ronner E, Boersma E, Akkerhuis KM, *et al*. Patients with acute coronary syndromes without persistent ST elevation undergoing percutaneous coronary intervention benefit most from early intervention with protection by a glycoprotein IIb/IIIa receptor blocker. *Eur Heart J* 2002;**23**:239–46.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2005.082826

Intramycocardial echinococcal cyst demonstrated by multislice computed tomography



A 52-year-old man was admitted to our hospital because of palpitations and fatigue. The physical examination was normal. A 12 lead ECG showed sinus tachycardia. The indirect haemagglutination test was negative but there was pronounced eosinophilia. Transthoracic echocardiography showed a suspicious lesion in the left ventricular apex, which could have been a cardiac hydatid cyst. Transoesophageal echocardiography was proposed but the patient was unable to tolerate it. Enhanced multislice computed tomography (MCT) was performed. Volume rendered images revealed a 3 × 2.5 cm cystic structure planed in left ventricular apex (panel A: C, cyst; D1, diagonal artery; LAD, left anterior descending artery). An axial maximal intensity projection image clearly showed septates within the cyst (panel B: LV, left ventricle; RV, right ventricle). With a diagnosis of echinococcosis, the patient was referred for surgical resection of the mass. Analysis of the resected specimen revealed it to be a hydatid cyst.

Transthoracic echocardiography is usually the initial diagnostic test in patients with a suspected cardiac mass. However, this technique is restricted by its small field of views and insufficient acoustic window in some patients. MCT allows for detailed delineation of intra- and pericardiac masses. As a result, we wanted to present the MCT images of the left ventricular cyst that we could not evaluate echocardiographically.

F Gundogdu
S Arslan
A M Kantarci
doktorfuat@hotmail.com